arXiv:1212.1059v1 [math.CA] 5 Dec 2012

On the degree of strong approximation of almost periodic functions in the Stepanov sense

Włodzimierz Łenski and Bogdan Szal
University of Zielona Góra
Faculty of Mathematics, Computer Science and Econometrics
65-516 Zielona Góra, ul. Szafrana 4a, Poland
W.Lenski@wmie.uz.zgora.pl, B.Szal @wmie.uz.zgora.pl

Abstract

Considering the class of almost periodic functions in the Stepanov sense we extend and generalize the results of the first author [11]. as well as the results of L. Leindler [7] and P. Chandra [4, 5] .

Key words: Degree of strong approximation, Almost periodic functions, Strong approximation, Special sequences.

2000 Mathematics Subject Classification: 42A24, 41A25.

1 Introduction

Let S^p $(1<p\leq\infty)$ be the class of all almost periodic functions in the Stepanov sense $(1<p<\infty)$ or uniformly almost periodic $(p=\infty)$ with the norm

$$\|f\|_{S^p}:=\begin{cases}\sup\limits_u\left\{\frac{1}{\pi}\int_u^{u+\pi}\mid f(t)\mid^p dt\right\}^{1/p} & \text{when}\quad 1<p<\infty,\\ \sup\limits_u\mid f(u)\mid \quad \text{when}\quad p=\infty.\end{cases}$$

Suppose that the Fourier series of $f\in S^p$ has the form

$$Sf(x)=\sum_{\nu=-\infty}^{\infty}A_\nu(f)\,e^{i\lambda_\nu x},\quad \text{where}\ \ A_\nu(f)=\lim_{L\to\infty}\frac{1}{L}\int_0^L f(t)e^{-i\lambda_\nu t}dt,$$

with the partial sums

$$S_{\gamma_k}f(x)=\sum_{|\lambda_\nu|\leq\gamma_k}A_\nu(f)\,e^{i\lambda_\nu x}$$

and that $0=\lambda_0<\lambda_\nu<\lambda_{\nu+1}$ if $\nu\in\mathbb{N}=\{1,2,3...\}$, $\lim\limits_{v\to\infty}\lambda_\nu=\infty$, $\lambda_{-\nu}=-\lambda_\nu$, $|A_\nu|+|A_{-\nu}|>0$. Let $\Omega_{\alpha,p}$, with some fixed positive α , be the set of functions of class S^p whose Fourier exponents satisfy the condition

$$\lambda_{\nu+1}-\lambda_\nu\ge\alpha\quad(\nu\in\mathbb{N}).$$

In case $f\in\Omega_{\alpha,p}$

$$S_{\lambda_k}f(x)=\int_0^\infty\{f(x+t)+f(x-t)\}\,\Psi_{\lambda_k,\lambda_k+\alpha}(t)\,dt,$$

where

$$\Psi_{\lambda,\eta}(t)=\frac{2\sin\frac{(\eta-\lambda)t}{2}\sin\frac{(\eta+\lambda)t}{2}}{\pi(\eta-\lambda)t^2}\qquad(0<\lambda<\eta,\quad|t|>0).$$

Let $A:=(a_{nk})$ $(k,n=0,1,...)$ be a lower triangular infinite matrix of real numbers satisfying the following condition:

$$a_{nk}\ge0\ \ (k,n=0,1,...),\ \ a_{nk}=0\ \ (k>n)\ \text{ and }\sum_{k=0}^n a_{nk}=1. \tag{1}$$

Let us consider the strong mean

$$H^q_{n,A,\gamma}f(x)=\left\{\sum_{k=0}^n a_{n,k}\left|S_{\gamma_k}f(x)-f(x)\right|^q\right\}^{1/q}\qquad(q>0). \tag{2}$$

As measures of approximation by the quantity (2), we use the best approximation of f by entire functions g_σ of exponential type σ bounded on the real axis, shortly $g_\sigma\in B_\sigma$ and the moduli of continuity of f defined by the formulas

$$E_\sigma(f)_{S^p}=\inf_{g_\sigma}\|f-g_\sigma\|_{S^p},$$

$$\omega f(\delta)_{S^p}=\sup_{|t|\le\delta}\|f(\cdot+t)-f(\cdot)\|_{S^p}$$

and

$$w_xf(\delta)_p:=\left\{\frac{1}{\delta}\int_0^\delta|\varphi_x(t)|^p\,dt\right\}^{1/p}\ \text{ with }1<p<\infty,$$

where $\varphi_x(t):=f(x+t)+f(x-t)-2f(x)$, respectively.

A sequence $c:=(c_n)$ of nonnegative numbers tending to zero is called the Rest Bounded Variation Sequence, or briefly $c\in RBVS$, if it has the property

$$\sum_{k=m}^\infty|c_n-c_{n+1}|\le K(c)\,c_m \tag{3}$$

for all natural numbers m, where $K(c)$ is a constant depending only on c.

A sequence $c := (c_n)$ of nonnegative numbers will be called the Head Bounded Variation Sequence, or briefly $c \in HBVS$, if it has the property

$$\sum_{k=0}^{m-1} |c_n - c_{n+1}| \leq K(c)\, c_m \tag{4}$$

for all natural numbers m, or only for all $m \leq N$ if the sequence c has only finite nonzero terms and the last nonzero terms is c_N.

Therefore we assume that the sequence $(K(\alpha_n))_{n=0}^{\infty}$ is bounded, that is, that there exists a constant K such that

$$0 \leq K(\alpha_n) \leq K$$

holds for all n, where $K(\alpha_n)$ denote the sequence of constants appearing in the inequalities (3) or (4) for the sequence $\alpha_n := (a_{nk})_{k=0}^{\infty}$.Now we can give the conditions to be used later on. We assume that for all n and $0 \leq m \leq n$

$$\sum_{k=m}^{\infty} |a_{nk} - a_{nk+1}| \leq K a_{nm} \tag{5}$$

and

$$\sum_{k=0}^{m-1} |a_{nk} - a_{nk+1}| \leq K a_{nm} \tag{6}$$

hold if $\alpha_n := (a_{nk})_{k=0}^{\infty}$ belongs to $RBVS$ or $HBVS$, respectively.

The C-norm of the deviation $\left|\sum_{k=0}^{n} a_{n,k}\left[S_k f(x) - f(x)\right]\right|$, with the partial sums $S_k f$ of classical trigonometric Fourier series, was estimated by P. Chandra [4] [5] for monotonic sequences (a_{nk}) and by L. Leindler [7] for the sequences of bounded variation. These results were generalized by W. Łenski [11] who considered the strong means $H_{n,A}^{q}$, also in classical case, and the functions belonging to the L^p. In present paper we shall considered the almost periodic functions from the Stepanov class giving similarly estimations for the strong means $H_{n,A}^{q}$ in individual points and in norms.

We shall write $I_1 \ll I_2$ if there exists a positive constant C such that $I_1 \leq C I_2$.

2 Main results

Let us consider a function w_x of modulus of continuity type on the interval $[0, +\infty)$, i.e. a nondecreasing continuous function having the following properties: $w_x(0) = 0$, $w_x(\delta_1 + \delta_2) \leq w_x(\delta_1) + w_x(\delta_2)$ for any $\delta_1, \delta_2 \geq 0$ with x such that the set

$$\Omega_{\alpha,p}(w_x) = \left\{ f \in \Omega_{\alpha,p} : \left[\frac{1}{\delta}\int_0^{\delta} |\varphi_x(t) - \varphi_x(t \pm \gamma)|^p \, dt\right]^{1/p} \ll w_x(\gamma) \right.$$
$$\left. \text{and} \quad w_x f(\delta)_p \ll w_x(\delta) \quad , \quad \text{where} \quad \gamma, \delta > 0 \right\}$$

is nonempty. It is clear that $\Omega_{\alpha,p}(w_x) \subseteq \Omega_{\alpha,p'}(w_x)$, for $p' \leq p < \infty$.

Our main results are the following:

Theorem 1 *Let (1) and (6) hold. Suppose w_x is such that*

$$\left\{ u^{\frac{p}{q}} \int\limits_{u}^{\pi} \frac{(w_x(t))^p}{t^{1+\frac{p}{q}}} dt \right\}^{\frac{1}{p}} = O(uH_x(u)) \quad as \quad u \to 0^+, \tag{7}$$

where $H_x(u) \geq 0$, $1 < p \leq q$ and

$$\int\limits_{0}^{t} H_x(u)\, du = O(tH_x(t)) \quad as \quad t \to 0^+. \tag{8}$$

If $f \in \Omega_{\alpha,p}(w_x)$, then

$$H^q_{n,A,\gamma} f(x) = O\left(a_{nn} H_x(a_{nn}) + \left\{ \sum_{k=0}^{n} a_{n,k} \left(E_{\alpha k/2}(f)_{S^p} \right)^q \right\}^{1/q} \right), \tag{9}$$

where q is such that $1 < q(q-1)^{-1} \leq p \leq q$.

Theorem 2 *Let (1), (5), (7) and (8) hold. If $f \in \Omega_{\alpha,p}(w_x)$, then*

$$H^q_{n,A,\gamma} f(x) = O\left(a_{n0} H_x(a_{n0}) + \left\{ \sum_{k=0}^{n} a_{n,k} \left(E_{\alpha k/2}(f)_{S^p} \right)^q \right\}^{1/q} \right), \tag{10}$$

where q is such that $1 < q(q-1)^{-1} \leq p \leq q$.

Consequently, we can immediately derive the results on norm approximation.

Theorem 3 *Let (1) and (6) hold. Suppose $\omega f(\cdot)_{S^{\widetilde{p}}}$ is such that*

$$\left\{ u^{\frac{p}{q}} \int\limits_{u}^{\pi} \frac{(\omega f(t)_{S^{\widetilde{p}}})^p}{t^{1+\frac{p}{q}}} dt \right\}^{\frac{1}{p}} = O(uH(u)) \quad as \quad u \to 0^+ \tag{11}$$

holds, with $1 < p \leq q \leq \widetilde{p}$, where additionally H (≥ 0) instead of H_x satisfies the condition (8). If $f \in \Omega_{\alpha,\widetilde{p}}$, then

$$\left\| H^{q'}_{n,A,\gamma} f(\cdot) \right\|_{S^{\widetilde{p}}} = O(a_{nn} H_x(a_{nn})),$$

with $q' \in (0,q]$, where q is such that $1 < q(q-1)^{-1} \leq p \leq q$.

Theorem 4 *Let (1) and (5) hold. Suppose* $\omega f(\cdot)_{S^{\widetilde{p}}}$ *is such that (11) holds, with* $1 < p \leq q \leq \widetilde{p}$ *, where additionally* H (≥ 0) *instead of* H_x *satisfies the condition (8). If* $f \in \Omega_{\alpha,\widetilde{p}}$*, then*

$$\left\| H^{q'}_{n,A,\gamma} f(\cdot) \right\|_{S^{\widetilde{p}}} = O\left(a_{n0} H_x\left(a_{n0}\right)\right),$$

with $q' \in (0, q]$*, where* q *is such that* $1 < q\left(q-1\right)^{-1} \leq p \leq q$*.*

Remark 1 *Analyzing our proofs and dividing the integral in the formula*

$$\left\{ \sum_{k=0}^{n} a_{n,k} \left| \int_{0}^{\infty} \varphi_x(t)\, \Psi_{k+\kappa}(t)\, dt \right|^{q} \right\}^{1/q}$$

into parts with $\frac{\pi}{n+1}$ *instead of* $a_{n,n}$ *or* $a_{n,0}$ *we can obtain the next series of theorems analogously as in [11].*

3 Lemmas

To prove our theorems we need the following lemmas.

Lemma 1 *[11] If (7) and (8) hold, then*

$$\int_{0}^{u} \frac{w_x f(t)}{t} dt = O\left(u H_x(u)\right) \quad \left(u \to 0_{+}\right). \tag{12}$$

Lemma 2 *[16, Theorem 5.20 II, Ch. XII] Suppose that* $1 < q\left(q-1\right)^{-1} \leq p \leq q$ *and* $\xi = \frac{1}{p} + \frac{1}{q} - 1$*. If* $\left|t^{-\xi} g(t)\right| \in L^{p}$*, then*

$$\left\{ \frac{\left|a_0(g)\right|^q}{2} + \sum_{k=0}^{\infty} \left(\left|a_k(g)\right|^q + \left|b_k(g)\right|^q \right) \right\}^{\frac{1}{q}} \ll \left\{ \int_{-\pi}^{\pi} \left|t^{-\xi} g(t)\right|^{p} dt \right\}^{\frac{1}{p}}. \tag{13}$$

4 Proofs of the Results

4.1 Proof of Theorem 1

In the proof we will use the following function $\Phi_x f(\delta,\nu) = \frac{1}{\delta} \int_{\nu}^{\nu+\delta} \varphi_x(u)\, du$, with $\delta = \delta_n = \frac{\pi}{n+1}$ and its estimate from [10, Lemma 1, p.218]

$$\left|\Phi_x f\left(\xi_1, \xi_2\right)\right| \leq w_x\left(\xi_1\right) + w_x\left(\xi_2\right) \tag{14}$$

for $f \in \Omega_{\alpha,p}\left(w_x\right)$ and any $\xi_1, \xi_2 > 0$.

Since, for $n = 0$ our estimate is evident we consider $n > 0$, only.

Denote by $S_k^* f$ the sums of the form

$$S_{\frac{\alpha k}{2}} f(x) = \sum_{|\lambda_\nu| \le \frac{\alpha k}{2}} A_\nu(f) e^{i\lambda_\nu x}$$

such that the interval $\left(\frac{\alpha k}{2}, \frac{\alpha(k+1)}{2}\right)$ does not contain any λ_ν. Applying Lemma 1.10.2 of [9] we easily verify that

$$S_k^* f(x) - f(x) = \int_0^\infty \varphi_x(t) \Psi_k(t)\, dt,$$

where $\varphi_x(t) := f(x+t) + f(x-t) - 2f(x)$ and $\Psi_k(t) = \Psi_{\frac{\alpha k}{2}, \frac{\alpha(k+1)}{2}}(t)$, i.e.

$$\Psi_k(t) = \frac{4 \sin \frac{\alpha t}{4} \sin \frac{\alpha(2k+1)t}{4}}{\alpha \pi t^2}$$

(see also [3], p.41). Evidently, if the interval $\left(\frac{\alpha k}{2}, \frac{\alpha(k+1)}{2}\right)$ contains a Fourier exponent λ_ν, then

$$S_{\frac{\alpha k}{2}} f(x) = S_{k+1}^* f(x) - \left(A_\nu(f) e^{i\lambda_\nu x} + A_{-\nu}(f) e^{-i\lambda_\nu x}\right).$$

Since (see [1, p.78] and [2, p. 7])

$$\left\{\sum_{\nu=-\infty}^{\infty} |A_\nu(f)|^q\right\}^{1/q} \le \|f\|_{B^p} \quad \text{and} \quad \|f\|_{B^p} \le \|f\|_{S^p},$$

where $\|\cdot\|_{B^p,}$ with $p \ge 1$, is the Besicovitch norm, so we have

$$|A_{\pm\nu}(f)| = \left|A_{\pm\nu}\left(f - g_{\alpha\mu/2}\right)\right| \le \left\|f - g_{\alpha\mu/2}\right\|_{S^p} = E_{\alpha\mu/2}(f)_{S^p},$$

for some $g_{\alpha\mu/2} \in B_{\alpha\mu/2}$, with $\alpha k/2 < \alpha\mu/2 < \lambda_\nu$. Therefore, the deviation

$$\left\{\sum_{k=0}^{n} a_{n,k} \left|S_{\frac{\alpha k}{2}} f(x) - f(x)\right|^q\right\}^{1/q}$$

can be estimated from above by

$$\left\{\sum_{k=0}^{n} a_{n,k} \left|\int_0^\infty \varphi_x(t) \Psi_{k+\kappa}(t)\, dt\right|^q\right\}^{1/q} + \left\{\sum_{k=0}^{n} a_{n,k} \left(E_{\alpha k/2}(f)_{S^p}\right)^q\right\}^{1/q},$$

where κ equals 0 or 1. Applying the Minkowski inequality we obtain

$$\left\{\sum_{k=0}^{n} a_{n,k} \left|\int_0^\infty \varphi_x(t) \Psi_{k+\kappa}(t)\, dt\right|^q\right\}^{1/q}$$

$$= \left\{ \sum_{k=0}^{n} a_{n,k} \left| \left(\int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} + \int_{\frac{2\pi}{\alpha}a_{n,n}}^{\frac{2\pi}{\alpha}} + \int_{\frac{2\pi}{\alpha}}^{\infty} \right) \varphi_x(t)\, \Psi_{k+\kappa}(t)\, dt \right|^q \right\}^{1/q}$$

$$\leq \left\{ \sum_{k=0}^{n} a_{n,k} \left| I_1(k) \right|^q \right\}^{1/q} + \left\{ \sum_{k=0}^{n} a_{n,k} \left| I_2(k) \right|^q \right\}^{1/q} + \left\{ \sum_{k=0}^{n} a_{n,k} \left| I_3(k) \right|^q \right\}^{1/q}.$$

By (1), integrating by parts, we obtain

$$\left\{ \sum_{k=0}^{n} a_{n,k} \left| I_1(k) \right|^q \right\}^{1/q} \leq \left\{ \sum_{k=0}^{n} a_{n,k} \left| \frac{4}{\alpha\pi} \int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} \varphi_x(t) \frac{\sin\frac{\alpha t}{4}}{t^2} \sin\frac{\alpha t}{4}(2k+2\kappa+1) dt \right|^q \right\}^{1/q}$$

$$\leq \frac{1}{\pi} \int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} \frac{|\varphi_x(t)|}{t} dt = \frac{1}{\pi} \int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} \frac{1}{t} \left(\frac{d}{dt} \int_{0}^{t} |\varphi_x(s)|\, ds \right) dt$$

$$= \frac{1}{\pi} \left[\frac{1}{t} \int_{0}^{t} |\varphi_x(s)|\, ds \right]_{t=0}^{t=\frac{2\pi}{\alpha}a_{n,n}} + \frac{1}{\pi} \int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} \frac{1}{t^2} \left(\int_{0}^{t} |\varphi_x(s)|\, ds \right) dt$$

$$= \frac{1}{\pi} w_x f\left(\frac{2\pi}{\alpha} a_{n,n} \right)_1 + \frac{1}{\pi} \int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} \frac{1}{t} w_x f(t)_1\, dt$$

$$\ll w_x f(a_{n,n})_1 + \int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} \frac{1}{t} w_x f(t)_1\, dt$$

$$= \frac{1}{\pi} w_x f\left(\frac{2\pi}{\alpha} a_{n,n} \right)_1 + \frac{1}{\pi} \int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} \frac{1}{t} w_x f(t)_1\, dt.$$

$$\ll w_x f(a_{n,n})_1 + \int_{0}^{\frac{2\pi}{\alpha}a_{n,n}} \frac{1}{t} w_x f(t)_1\, dt. \tag{15}$$

It is clear that $w_x f(\delta)_1 / \delta$ is nondecreasing with respect to $\delta > 0$ and $w_x f(\delta)_1 \leq w_x f(\delta)_p$ for $p \geq 1$. Using these properties we have

$$\left\{ \sum_{k=0}^{n} a_{n,k} \left| I_1(k) \right|^q \right\}^{1/q} \ll a_{n,n} \int_{a_{n,n}}^{\pi} \frac{w_x f(t)_1}{t^2} + \int_{0}^{a_{n,n}} \frac{1}{t} w_x f\left(\frac{2\pi}{\alpha} t \right)_1 dt$$

$$\ll \left\{ a_{n,n} \int\limits_{a_{n,n}}^{\pi} \frac{\left(w_x f\left(t\right)_1\right)^p}{t^2} \right\}^{\frac{1}{p}} + \int\limits_{0}^{a_{n,n}} \frac{1}{t} w_x f\left(t\right)_1 dt.$$

Since $f \in \Omega_{\alpha,p}\left(w_x\right)$ and (8) holds, Lemma 1 and (7) give

$$\left\{ \sum_{k=0}^{n} a_{n,k} \left|I_1(k)\right|^q \right\}^{1/q} = O\left(a_{nn} H_x\left(a_{nn}\right)\right).$$

If (6) holds, then

$$a_{n,\mu} - a_{n,m} \le \left|a_{n,\mu} - a_{n,m}\right| \le \sum_{k=\mu}^{m-1} \left|a_{n,k} - a_{n,k+1}\right| \le K a_{n,m}$$

for any $m \ge \mu \ge 0$. Hence we have

$$a_{n,\mu} \le \left(K+1\right) a_{n,m}. \tag{16}$$

From this, we get

$$\begin{aligned} & \left\{ \sum_{k=0}^{n} a_{n,k} \left|I_2(k)\right|^q \right\}^{1/q} \\ \le \quad & \left\{\left(K+1\right) a_{n,n}\right\}^{\frac{1}{q}} \left\{ \sum_{k=0}^{n} \left| \frac{4}{\alpha\pi} \int\limits_{\frac{2\pi}{\alpha} a_{n,n}}^{\frac{2\pi}{\alpha}} \frac{\varphi_x\left(t\right) \sin\frac{\alpha t}{4}}{t^2} \sin\frac{\alpha t}{4}(2k+2\kappa+1) dt \right|^q \right\}^{\frac{1}{q}} \\ \ll & \frac{8}{\alpha^2}\left(a_{nn}\right)^{\frac{1}{q}} \left\{ \sum_{k=0}^{n} \left| \frac{\alpha}{2\pi} \int\limits_{\frac{2\pi}{\alpha} a_{n,n}}^{\frac{2\pi}{\alpha}} \frac{\varphi_x\left(t\right) \sin\frac{\alpha t}{4}}{t^2} \sin\frac{\alpha t}{4}(2\kappa+1) \cos\frac{\alpha k t}{2} dt \right|^q \right\}^{\frac{1}{q}} \\ & + \frac{8}{\alpha^2}\left(a_{nn}\right)^{\frac{1}{q}} \left\{ \sum_{k=0}^{n} \left| \frac{\alpha}{2\pi} \int\limits_{\frac{2\pi}{\alpha} a_{n,n}}^{\frac{2\pi}{\alpha}} \frac{\varphi_x\left(t\right) \sin\frac{\alpha t}{4}}{t^2} \cos\frac{\alpha t}{4}(2\kappa+1) \sin\frac{\alpha k t}{2} dt \right|^q \right\}^{\frac{1}{q}}. \end{aligned}$$

Using inequality (13), we have

$$\left\{ \sum_{k=0}^{n} a_{n,k} \left|I_2(k)\right|^q \right\}^{1/q} \ll \left(a_{nn}\right)^{\frac{1}{q}} \left\{ \int\limits_{\frac{2\pi}{\alpha} a_{n,n}}^{\frac{2\pi}{\alpha}} \frac{\left|\varphi_x\left(t\right)\right|^p}{t^{1+\frac{p}{q}}} \right\}^{\frac{1}{p}}.$$

Integrating by parts, we obtain

$$\left\{ \sum_{k=0}^{n} a_{n,k} \left|I_2(k)\right|^q \right\}^{1/q} = \left(a_{nn}\right)^{\frac{1}{q}} \left\{ \left[\frac{1}{t^{1+\frac{p}{q}}} \int\limits_{0}^{t} \left|\varphi_x\left(t\right)\right|^p ds \right]_{t=\frac{2\pi}{\alpha} a_{n,n}}^{t=\frac{2\pi}{\alpha}} \right.$$

$$\left(1+\frac{p}{q}\right)\int\limits_{\frac{2\pi}{\alpha}a_{n,n}}^{\frac{2\pi}{\alpha}}\frac{1}{t^{2+\frac{p}{q}}}\left(\int\limits_{0}^{t}\left|\varphi_x\left(t\right)\right|^p ds\right)dt\Bigg\}^{\frac{1}{p}}$$

$$=\left(a_{nn}\right)^{\frac{1}{q}}\left\{\left[\frac{1}{t^{\frac{p}{q}}}\left(w_x f\left(t\right)_p\right)^p\right]_{t=\frac{2\pi}{\alpha}a_{n,n}}^{t=\frac{2\pi}{\alpha}}+\left(1+\frac{p}{q}\right)\int\limits_{\frac{2\pi}{\alpha}a_{n,n}}^{\frac{2\pi}{\alpha}}\frac{1}{t^{1+\frac{p}{q}}}\left(w_x f\left(t\right)_p\right)^p dt\right\}^{\frac{1}{p}}$$

$$\ll\left(a_{nn}\right)^{\frac{1}{q}}\left\{\left(w_x f\left(\frac{2\pi}{\alpha}\right)_p\right)^p+\int\limits_{\frac{2\pi}{\alpha}a_{n,n}}^{\frac{2\pi}{\alpha}}\frac{1}{t^{1+\frac{p}{q}}}\left(w_x f\left(t\right)_p\right)^p dt\right\}^{\frac{1}{p}}. \tag{17}$$

Since $f\in\Omega_{\alpha,p}\left(w_x\right)$, (7) gives

$$\left\{\sum_{k=0}^{n}a_{n,k}\left|I_2(k)\right|^q\right\}^{1/q}\ll\left(a_{nn}\right)^{\frac{1}{q}}\left\{\left(w_x\left(\pi\right)\right)^p+\int\limits_{a_{nn}}^{\pi}\frac{1}{t^{1+\frac{p}{q}}}\left(w_x\left(t\right)\right)^p dt\right\}^{\frac{1}{p}}$$

$$\ll\left\{\left(a_{nn}\right)^{\frac{p}{q}}\int\limits_{a_{nn}}^{\pi}\frac{\left(w_x\left(t\right)\right)^p}{t^{1+\frac{p}{q}}}dt\right\}^{\frac{1}{p}}=O\left(a_{nn}H_x\left(a_{nn}\right)\right).$$

For the third term we obtain

$$\left\{\sum_{k=0}^{n}a_{n,k}\left|I_3(k)\right|^q\right\}^{1/q}\le$$

$$\le\left\{\sum_{k=0}^{n}a_{n,k}\left|\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\left[\varphi_x\left(t\right)-\Phi_x f\left(\delta_k,t\right)\right]\Psi_{k+\kappa}\left(t\right)dt\right|^q\right\}^{1/q}$$

$$+\left\{\sum_{k=0}^{n}a_{n,k}\left|\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\Phi_x f\left(\delta_k,t\right)\Psi_{k+\kappa}\left(t\right)dt\right|^q\right\}^{1/q}$$

$$=\left\{\sum_{k=0}^{n}a_{n,k}\left|I_{31}(k)\right|^q\right\}^{1/q}+\left\{\sum_{k=0}^{n}a_{n,k}\left|I_{32}(k)\right|^q\right\}^{1/q}$$

and

$$\left|I_{31}(k)\right|\le\frac{4}{\alpha\pi}\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\left|\varphi_x\left(t\right)-\Phi_x f\left(\delta_k,t\right)\right|t^{-2}dt$$

$$\le \frac{4}{\alpha\pi}\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\left[\frac{1}{\delta_k t^2}\int_0^{\delta_k}\left|\varphi_x(t)-\varphi_x(t+u)\right|du\right]dt$$

$$=\frac{4}{\alpha\pi}\frac{1}{\delta_k}\int\limits_0^{\delta_k}\sum_{\mu=1}^{\infty}\left\{\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\frac{1}{t^2}\left|\varphi_x(t)-\varphi_x(t+u)\right|dt\right\}du$$

$$\begin{aligned}=\frac{4}{\alpha\pi}\frac{1}{\delta_k}\int\limits_0^{\delta_k}\sum_{\mu=1}^{\infty}\Bigg\{&\left[\frac{1}{t^2}\int_0^t\left|\varphi_x(s)-\varphi_x(s+u)\right|ds\right]_{t=\frac{2\pi}{\alpha}\mu}^{t=\frac{2\pi}{\alpha}(\mu+1)}\\&+2\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\left[\frac{1}{t^3}\int_0^t\left|\varphi_x(s)-\varphi_x(s+u)\right|ds\right]dt\Bigg\}du\end{aligned}$$

$$\begin{aligned}\ll\Bigg|\frac{1}{\delta_k}\int\limits_0^{\delta_k}\sum_{\mu=1}^{\infty}\Bigg\{&\frac{1}{[\frac{2\pi}{\alpha}(\mu+1)]^2}\int\limits_0^{\frac{2\pi}{\alpha}(\mu+1)}\left|\varphi_x(s)-\varphi_x(s+u)\right|ds\\&-\frac{1}{[\frac{2\pi}{\alpha}\mu]^2}\int\limits_0^{\frac{2\pi}{\alpha}\mu}\left|\varphi_x(s)-\varphi_x(s+u)\right|ds\Bigg\}du\Bigg|\\+\frac{1}{\delta_k}\int\limits_0^{\delta_k}\sum_{\mu=1}^{\infty}\Bigg\{&\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\left[\frac{1}{t^3}\int_0^t\left|\varphi_x(s)-\varphi_x(s+u)\right|ds\right]dt\Bigg\}du.\end{aligned}$$

Since $f\in\Omega_{\alpha,p}(w_x)$, thus for any x

$$\begin{aligned}\lim_{\zeta\to\infty}\frac{1}{\zeta^2}\int\limits_0^{\zeta}\left|\varphi_x(s)-\varphi_x(s+u)\right|ds &\le \lim_{\zeta\to\infty}\frac{1}{\zeta}w_x(u)\le\lim_{\zeta\to\infty}\frac{1}{\zeta}w_x(\delta_k)\\&\le\lim_{\zeta\to\infty}\frac{1}{\zeta}w_x(\pi)=0,\end{aligned}$$

and therefore

$$\begin{aligned}|I_{31}(k)|\le\frac{1}{\delta_k}\int\limits_0^{\delta_k}\frac{\alpha}{2\pi}\left[\frac{\alpha}{2\pi}\int_0^{2\pi/\alpha}\left|\varphi_x(s)-\varphi_x(s+u)\right|ds\right]du\\+\frac{1}{\delta_k}\int\limits_0^{\delta_k}w_x(u)\,du\sum_{\mu=1}^{\infty}\left\{\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\frac{1}{t^2}dt\right\}\end{aligned}$$

$$\ll \frac{1}{\delta_k}\int\limits_0^{\delta_k} w_x\left(u\right)du + w_x\left(\delta_k\right)\sum_{\mu=1}^{\infty}\frac{1}{\frac{2\pi}{\alpha}\mu^2} \ll w_x\left(\delta_k\right).$$

Next, we will estimate the term $\left|I_{32}(k)\right|$. So,

$$I_{32}(k)=\frac{2}{\alpha\pi}\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\frac{\Phi_x f\left(\delta_k,t\right)}{t^2}\frac{d}{dt}\left(-\frac{\cos\frac{\alpha t(k+\kappa)}{2}}{\frac{\alpha(k+\kappa)}{2}}+\frac{\cos\frac{\alpha t(k+\kappa+1)}{2}}{\frac{\alpha(k+\kappa+1)}{2}}\right)dt$$

$$=\frac{2}{\alpha\pi}\sum_{\mu=1}^{\infty}\left[\frac{\Phi_x f\left(\delta_k,t\right)}{t^2}\left(-\frac{\cos\frac{\alpha t(k+\kappa)}{2}}{\frac{\alpha(k+\kappa)}{2}}+\frac{\cos\frac{\alpha t(k+\kappa+1)}{2}}{\frac{\alpha(k+\kappa+1)}{2}}\right)\right]_{t=\frac{2\pi}{\alpha}\mu}^{t=\frac{2\pi}{\alpha}(\mu+1)}$$

$$+\frac{2}{\alpha\pi}\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\frac{d}{dt}\left(\frac{\Phi_x f\left(\delta_k,t\right)}{t^2}\right)\left(\frac{\cos\frac{\alpha t(k+\kappa)}{2}}{\frac{\alpha(k+\kappa)}{2}}-\frac{\cos\frac{\alpha t(k+\kappa+1)}{2}}{\frac{\alpha(k+\kappa+1)}{2}}\right)dt$$

$$= \quad I_{321}\left(k\right)+I_{322}\left(k\right)$$

Since $f\in\Omega_{\alpha,p}\left(w_x\right)$, thus for any x (using (14))

$$\lim_{\zeta\to\infty}\left|\frac{\Phi_x f\left(\delta_k,\frac{2\pi}{\alpha}\zeta\right)}{\left[\frac{2\pi}{\alpha}\zeta\right]^2}\left(-\frac{\cos\left[\pi\zeta(k+\kappa)\right]}{\frac{\alpha(k+\kappa)}{2}}+\frac{\cos\left[\pi\zeta\left(k+\kappa+1\right)\right]}{\frac{\alpha(k+\kappa+1)}{2}}\right)\right|$$

$$\leq\lim_{\zeta\to\infty}\frac{w_x\left(\delta_k\right)+w_x\left(\frac{2\pi}{\alpha}\zeta\right)}{2\pi^2\zeta^2 k}\ll\lim_{\zeta\to\infty}\frac{w_x\left(\delta_k\right)+\zeta w_x\left(\frac{2\pi}{\alpha}\right)}{\zeta^2 k}\ll w_x\left(\pi\right)\lim_{\zeta\to\infty}\frac{1+\zeta}{\zeta^2}=0,$$

and therefore

$$I_{321}\left(k\right)=\frac{2}{\alpha\pi}\sum_{\mu=1}^{\infty}\left[\frac{\Phi_x f\left(\delta_k,\frac{2\pi}{\alpha}\left(\mu+1\right)\right)}{\left[\frac{2\pi}{\alpha}\left(\mu+1\right)\right]^2}\left(-\frac{\cos\left[\pi\left(\mu+1\right)\left(k+\kappa\right)\right]}{\frac{\alpha(k+\kappa)}{2}}\right.\right.$$

$$\left.+\frac{\cos\left[\pi\left(\mu+1\right)\left(k+\kappa+1\right)\right]}{\frac{\alpha(k+\kappa+1)}{2}}\right)$$

$$\left.-\frac{\Phi_x f\left(\delta_k,\frac{2\pi}{\alpha}\mu\right)}{\left[\frac{2\pi}{\alpha}\mu\right]^2}\left(-\frac{\cos\left[\pi\mu(k+\kappa)\right]}{\frac{\alpha(k+\kappa)}{2}}+\frac{\cos\left[\pi\mu\left(k+\kappa+1\right)\right]}{\frac{\alpha(k+\kappa+1)}{2}}\right)\right]$$

$$=-\frac{2}{\alpha\pi}\frac{\Phi_x f\left(\delta_k,2\pi/\alpha\right)}{\left[2\pi/\alpha\right]^2}\left(-\frac{\left(-1\right)^{(k+\kappa)}}{\frac{\alpha(k+\kappa)}{2}}+\frac{\left(-1\right)^{(k+\kappa+1)}}{\frac{\alpha(k+\kappa+1)}{2}}\right)$$

$$=-\frac{1}{\pi^3}\Phi_x f\left(\delta_k,2\pi/\alpha\right)\left(-1\right)^{(k+\kappa+1)}\left(\frac{1}{k+\kappa+1}+\frac{1}{k+\kappa}\right).$$

Using (14), we get

$$|I_{321}(k)| \ll \frac{1}{\pi^3}\frac{2}{k+1}\left|\Phi_x f\left(\delta_k, 2\pi/\alpha\right)\right| \le \frac{2}{\pi^3(k+1)}\left(w_x(\delta_k) + w_x(2\pi/\alpha)\right).$$

Similarly

$$I_{322}(k) = \frac{2}{\alpha\pi}\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\left(\frac{\frac{d}{dt}\Phi_x f(\delta_k,t)}{t^2} - \frac{2\Phi_x f(\delta_k,t)}{t^3}\right)$$

$$\cdot\left(\frac{\cos\frac{\alpha t(k+\kappa)}{2}}{\frac{\alpha(k+\kappa)}{2}} - \frac{\cos\frac{\alpha t(k+\kappa+1)}{2}}{\frac{\alpha(k+\kappa+1)}{2}}\right)dt$$

and

$$|I_{322}(k)| \ll \frac{8}{\alpha^2(k+1)\pi}\sum_{\mu=1}^{\infty}\left[\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\frac{|\varphi_x(t+\delta_k) - \varphi_x(t)|}{\delta_k t^2}dt\right.$$

$$\left.+\,2\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\frac{|\Phi_x f(\delta_k,t)|}{t^3}dt\right]$$

$$\le \frac{8}{\alpha^2(k+1)\pi\delta_k}\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\frac{|\varphi_x(t+\delta_k) - \varphi_x(t)|}{t^2}dt$$

$$+\frac{16}{\alpha^2(k+1)\pi}\sum_{\mu=1}^{\infty}\int\limits_{\frac{2\pi}{\alpha}\mu}^{\frac{2\pi}{\alpha}(\mu+1)}\frac{w_x(\delta_k) + w_x(t)}{t^3}dt$$

$$\ll \frac{1}{(k+1)\delta_k}w_x(\delta_k) + \frac{1}{k+1}\sum_{\mu=1}^{\infty}\left[\left(w_x(\delta_k) + w_x\left(\frac{2\pi(\mu+1)}{\alpha}\right)\right)\frac{\alpha^2}{4\pi^2\mu^3}\right]$$

$$\ll w_x(\delta_k) + \frac{1}{k+1}\left[w_x(\delta_k)\sum_{\mu=1}^{\infty}\frac{1}{\mu^3} + \sum_{\mu=1}^{\infty}\frac{w_x\left(\frac{2\pi(\mu+1)}{\alpha}\right)}{\mu^3}\right]$$

$$\ll w_x(\delta_k) + \frac{1}{k+1}\left(w_x(\delta_k) + w_x\left(\frac{4\pi}{\alpha}\right)\sum_{\mu=1}^{\infty}\frac{\mu+1}{\mu^3}\right)$$

$$\ll w_x(\delta_k) + \frac{1}{k+1}\left(w_x(\delta_k) + w_x\left(\frac{4\pi}{\alpha}\right)\right).$$

Therefore

$$|I_3(k)| \ll w_x(\delta_k) + \frac{1}{k+1}\left(w_x(\delta_k) + w_x\left(\frac{2\pi}{\alpha}\right) + w_x\left(\frac{4\pi}{\alpha}\right)\right)$$

and thus

$$\left\{\sum_{k=0}^{n} a_{n,k}\left|I_3(k)\right|^q\right\}^{1/q} \ll \left\{\sum_{k=0}^{n} a_{n,k}\left(w_x\left(\frac{\pi}{k+1}\right)+\frac{1}{k+1}w_x\left(\frac{\pi}{\alpha}\right)\right)^q\right\}^{1/q}$$

$$\ll \left\{\sum_{k=0}^{n} a_{n,k}\left(w_x\left(\frac{\pi}{k+1}\right)\right)^q\right\}^{1/q}.$$

From (16) we obtain

$$\sum_{k=0}^{n} a_{n,k}\left(w_x\left(\frac{\pi}{k+1}\right)\right)^q \leq \sum_{k=0}^{\left[\frac{1}{(K+1)a_{n,n}}\right]-1} a_{nk}\left(w_x\left(\frac{\pi}{k+1}\right)\right)^q$$

$$+\sum_{k=\left[\frac{1}{(K+1)a_{n,n}}\right]-1}^{n} a_{nk}\left(w_x\left(\frac{\pi}{k+1}\right)\right)^q.$$

Using (1), (16) and the monotonicity of the function w_x, from (7) and (12), we get

$$\sum_{k=0}^{n} a_{n,k}\left(w_x\left(\frac{\pi}{k+1}\right)\right)^q \leq (K+1)\,a_{n,n}\sum_{k=0}^{\left[\frac{1}{(K+1)a_{n,n}}\right]-1}\left(w_x\left(\frac{\pi}{k+1}\right)\right)^q$$

$$+\left(w_x\left(\pi\left(K+1\right)a_{n,n}\right)\right)^q\sum_{k=\left[\frac{1}{(K+1)a_{n,n}}\right]-1}^{n} a_{nk}$$

$$\ll a_{n,n}\int_{1}^{\frac{1}{(K+1)a_{n,n}}}\left(w_x\left(\frac{\pi}{t}\right)\right)^q dt+\left(w_x\left(a_{n,n}\right)\right)^q \ll a_{n,n}\int_{a_{n,n}}^{\pi}\frac{\left(w_x\left(u\right)\right)^q}{u^2}du+\left(w_x\left(a_{n,n}\right)\right)^q$$

$$\leq a_{n,n}\int_{a_{n,n}}^{\pi}\frac{\left(w_x\left(u\right)\right)^q}{u^{1+p/q+1-p/q}}du+\left(4w_x\left(\frac{a_{n,n}}{2}\right)\right)^q$$

$$\leq \left(a_{n,n}\right)^{p/q}\int_{a_{n,n}}^{\pi}\frac{\left(w_x\left(u\right)\right)^q}{u^{1+p/q}}du+\left(8\int_{\frac{a_{n,n}}{2}}^{a_{n,n}}\frac{w_x\left(u\right)}{u}du\right)^q$$

$$\ll \left(a_{n,n}\right)^{p/q}\int_{a_{n,n}}^{\pi}\frac{\left(w_x\left(u\right)\right)^q}{u^{1+p/q}}du+\left(\int_{0}^{a_{n,n}}\frac{w_x\left(u\right)}{u}du\right)^q \ll \left(a_{n,n}H_x\left(a_{n,n}\right)\right)^q.$$

Summing up we obtain that (9) is proved and the proof is complete.

4.2 Proof of Theorem 2

Under the notation of the before proof we can write

$$\left\{\sum_{k=0}^{n} a_{n,k}\left|\int_{0}^{\infty}\varphi_x\left(t\right)\Psi_{k+\kappa}\left(t\right)dt\right|^{q}\right\}^{1/q}$$

$$=\left\{\sum_{k=0}^{n} a_{n,k}\left|\left(\int\limits_{0}^{\frac{2\pi}{\alpha}a_{n,0}}+\int\limits_{\frac{2\pi}{\alpha}a_{n,0}}^{\frac{2\pi}{\alpha}}+\int\limits_{\frac{2\pi}{\alpha}}^{\infty}\right)\varphi_x\left(t\right)\Psi_{k+\kappa}\left(t\right)dt\right|^{q}\right\}^{1/q}$$

$$\leq\left\{\sum_{k=0}^{n} a_{n,k}\left|J_1(k)\right|^{q}\right\}^{1/q}+\left\{\sum_{k=0}^{n} a_{n,k}\left|J_2(k)\right|^{q}\right\}^{1/q}+\left\{\sum_{k=0}^{n} a_{n,k}\left|J_3(k)\right|^{q}\right\}^{1/q},$$

using the Minkowski inequality. Applying the property of the class $RBVS$ instead of the property of $HBVS$ our proof will be similar to the proof of Theorem 1.

References

[1] A. Avantaggiati, G. Bruno and B. Iannacci, The Hausdorff-Young theorem for almost periodic functions and some applications, Nonlinear analysis, Theory, Methods and Applications, Vol. 25, No. 1, (1995), pp. 61-87.

[2] A. D. Bailey, Almost Everywhere Convergence of Dyadic Partial Sums of Fourier Series for Almost Periodic Functions, Master of Philosophy, A thesis submitted to School of Mathematics of The University of Birmingham for the degree of Master of Philosophy, September 2008.

[3] A. S. Besicovitch, Almost periodic functions, Cambridge, 1932.

[4] P. Chandra, On the degree of approximation of a class of functions by means of Fourier series, Acta Mat. Hung., 52 (1988), 199-205.

[5] P. Chandra, A note on the degree of approximation of continuous functions, Acta Math. Hungar., 62 (1993), 21-23.

[6] L. Leindler, On the degree of approximation of continuous functions, Acta Math. Hungar., 104 (2004), 105-113.

[7] L Leindler, On the uniform convergence and boundedness of a certain class of sine series, Analysis Math., 27 (2001), 279-285.

[8] L. Leindler, A new extension of monotone sequence and its application, J. Inequal. Pure and Appl. Math., 7(1) (2006), Art. 39, 7 pp.

[9] B. L. Levitan, Almost periodic functions, Gos. Izdat. Tekh-Teoret. Liter., Moscov 1953 (in Russian).

[10] W. Łenski, Pointwise strong and very strong approximation of Fourier series, Acta Math. Hung., 115(3), (2007), p.215-233.

[11] W. Łenski, On the degree of strong approximation of integrable functions, J. Inequal. Pure and Appl. Math., 8(3) (2007), Art. 70.

[12] P. Pych-Taberska, Approximation properties of the partial sums of Fourier series of almost periodic functions, Studia Math. XCVI (1990),91-103.

[13] S. Tikhonov, Trigonometric series with general monotone coefficients, J. Math. Anal. Appl., 326(1) (2007), 721-735.

[14] S. Tikhonov, On uniform convergence of trigonometric series. Mat. Zametki, 81(2) (2007), 304-310, translation in Math. Notes, 81(2) (2007), 268-274.

[15] S. Tikhonov, Best approximation and moduli of smoothness: Computation and equivalence theorems, J. Approx. Theory, 153 (2008), 19-39.

[16] A. Zygmund, Trigonometric Series, Cambridge, 2002.